arXiv:1512.04585v1 [math.FA] 11 Dec 2015

[1]

AN INEQUALITY FOR t-GEOMETRIC MEANS

TRUNG HOA DINH[A]

Abstract. In this note we prove an inequality for t-geometric means that immediately implies the recent results of Audenaert [2] and Hayajneh-Kittaneh [6].

1. Introduction

Recall that a norm $|||\cdot|||$ on the algebra $\mathbb{C}_{n\times n}$ of $n\times n$ complex matrices is unitarily invariant if $|||UAV||| = |||A|||$ for any unitary matrices $U, V \in \mathbb{C}_{n\times n}$ and any $A \in \mathbb{C}_{n\times n}$.

In [2] Audenaert proved that if $A_i, B_i \in \mathbb{P}_n$ $(i = 1, \cdots, m)$, such that $A_iB_i = B_iA_i$, then for any unitarily invariant norm $|||\cdot|||$ on $\mathbb{C}_{n\times n}$,

$$|||\sum_{i=1}^m A_iB_i||| \leqslant |||(\sum_{i=1}^m A_i^{1/2}B_i^{1/2})^2||| \leqslant |||(\sum_{i=1}^m A_i)(\sum_{i=1}^m B_i)|||\,. \tag{1.1}$$

In particular, this result confirms a conjecture of Hayajneh and Kittaneh in [6] and answers a question of Bourin. Very recently Lin [7] gave another proof of inequality (1.1). In the next section, based on a result of Bourin and Uchiyama in [4] we prove an inequality for t-geometric means that immediately implies (1.1).

2. Main result

The following lemma is follows from the well-known Hiai-Ando log-majorization Theorem and [8, Proposition 2.2].

Lemma 2.1. Let $A, B \in \mathbb{P}_n$ and $t \in [0, 1]$ and $s > 0$. For all unitarily invariant norms $|||\cdot|||$ on $\mathbb{C}_{n\times n}$,

$$|||(A\sharp_t B)^r||| \leqslant |||A^r\sharp_t B^r||| \leqslant |||\left(B^{rts/2}A^{(1-t)rs}B^{rts/2}\right)^{1/s}||| \leqslant |||\left(A^{(1-t)rs}B^{rts}\right)^{1/s}|||\,.$$

The following theorem was obtained by Bourin and Uchiyama [4, Theorem 1.2].

Theorem 2.2. *Let $A_i \geqslant 0$ $(i = 1, \cdots, m)$. Then for every non-negative convex function f on $[0, \infty)$ with $f(0) = 0$ and for any unitarily invariant norm $|||\cdot|||$*

$$|||f(A_1) + f(A_2) + \cdots f(A_m)||| \leqslant |||f(A_1 + A_2 + \cdots + A_m)|||. \tag{2.1}$$

The above inequality is reversed if f is non-negative concave function on $[0, \infty)$ with $f(0) = 0$.

Now we will prove our main theorem.

[1A]Research of the first author is funded by Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grant number 101.04-2014.40.

Date: December 16, 2015.

2010 *Mathematics Subject Classification.* 15A45, 15B48, 53C35.

Key words and phrases. t-geometric mean, positive definite matrices, log-majorization, unitarily invariant norms.

Theorem 2.3. Let $A_i, B_i \in M_n^+$, $i = 1, \ldots, m$. Then for any unitarily invariant norm $|||\cdot|||$ on M_n and $r \geqslant 1$ we have

$$|||\sum_{i=1}^{m}(A_i\sharp_t B_i)^r||| \leqslant |||(\sum_{i=1}^{m}A_i)^{r/4}(\sum_{i=1}^{m}B_i)^{r/2}(\sum_{i=1}^{m}A_i)^{r/4}||| \leqslant |||(\sum_{i=1}^{m}A_i)^{r/2}(\sum_{i=1}^{m}B_i)^{r/2}|||. \tag{2.2}$$

Proof. Since the function x^r is convex and monotone increasing, by Theorem 2.2 and the concavity of the t-geometric means, we have

$$|||\sum_{i=1}^{m}(A_i\sharp_t B_i)^r||| \leqslant |||(\sum_{i=1}^{m}A_i\sharp_t B_i)^r||| \leqslant |||\big((\sum_{i=1}^{m}A_i)\sharp_t(\sum_{i=1}^{m}B_i)\big)^r|||. \tag{2.3}$$

On account of Lemma 2.1

$$|||\big((\sum_{i=1}^{m}A_i)\sharp_t(\sum_{i=1}^{m}B_i)\big)^r||| \leqslant |||(\sum_{i=1}^{m}A_i)^{r/4}(\sum_{i=1}^{m}B_i)^{r/2}(\sum_{i=1}^{m}A_i)^{r/4}||| \leqslant |||(\sum_{i=1}^{m}A_i)^{r/2}(\sum_{i=1}^{m}B_i)^{r/2}|||. \tag{2.4}$$

Combining (2.3) and (2.4), we get (2.2). □

Remark 2.4. The case when $r = 2$ was considered in [8]. The results of Audenaert [2] and Hayajneh-Kittaneh [6] are the special case of (2.2) when $r = 2$ and $t = \frac{1}{2}$.

References

[1] T. Ando, F. Hiai, Log majorization and complementary Golden-Thompson type inequalities, Linear Algebra Appl., **197/198** (1994), 113–131.

[2] K.M.R. Audenaert, A norm inequality for pairs of commuting positive semidefinite matrices, Electron. J. Linear Algebra, **30** (2015), 80–84.

[3] R. Bhatia, *Matrix Analysis*, Springer, New York, 1997.

[4] J.C. Bourin, M. Uchiyama, A matrix subadditivity inequality for $f(A + B)$ and $f(A) + f(B)$, Linear Algebra Appl., **423** (2007), 512–518.

[5] J.C. Bourin, Matrix subadditivity inequalities and block-matrices, Internat. J. Math. **20** (2009), 679–691.

[6] S. Hayajneh, F. Kittaneh, Trace inequalities and a question of Bourin, Bull. Aust. Math. Soc. **88** (2013), 384–389.

[7] M. Lin, Remarks on two recent results of Audenaert, Linear Algebra Appl., **489** (2016), 24–29.

[8] Dinh Trung Hoa, Tin-Yau Tam, Geometry and Inequalities of Geometric Mean. Submitted, 2015.

Division of Computational Mathematics and Engineering, Institute for Computational Science, Ton Duc Thang University, Ho Chi Minh City, Vietnam, Faculty of Civil Engineering, Ton Duc Thang University, Vietnam, Department of Mathematics and Statistics, Auburn University, Auburn, AL 36849, USA,

E-mail address: dinhtrunghoa@tdt.edu.vn, trunghoa.math@gmail.com